



Guidance for Schools: Common Signs & Symptoms



This document is intended to support decision-making for schools and childcare centers when an individual child is displaying signs and symptoms of illness.

Below are suggestions for when to keep your child home when they are not feeling well. Contact a health care provider if your child needs medical care. A doctor’s note is generally not required to return to school or child care. For more information, please visit [the CDPH guidance page](#).

This document:

- **does not** replace the advice of a licensed healthcare provider, nor does it revise guidance related to an individual child’s known medical condition or existing IEP, Section 504 plan, or IFSP plan.
- **is not** a diagnostic tool. Related symptoms may inform exclusion decisions, but school staff should not attempt to diagnose.
- **does not** replace or revise applicable laws or other requirements.
- **does not** address specific disease outbreaks, when recommendations and public health requirements for exclusion may be more strict.
- **is limited to** consideration of symptoms as evidence of infection, and does not include considerations for other causes of symptoms such as allergies, toxin exposure, and child abuse or neglect.

Symptom of Illness	When should my child stay home and when can they return?
Overall, not feeling well	<p>STAY HOME IF any new illness or symptom prevents a child from participating meaningfully in routine activities.</p> <p>Please note that a child with multiple symptoms is more likely to be contagious. Recommendations for all applicable symptoms should be considered.</p>
Fever	<p>STAY HOME IF a fever is 100.4°F (38°C) or higher.</p> <p>RETURN WHEN fever went away over the night and is gone in the morning without the use of medications such as Tylenol[®], Advil[®], or Motrin[®] (acetaminophen or ibuprofen).</p>
Respiratory Virus Symptoms (cough, stuffy / runny nose, sneeze)	<p>OK TO ATTEND WITH MILD SYMPTOMS. <u>Children should test for COVID-19 before returning to school.</u> Those who test negative using an antigen test but continue having COVID-19 symptoms should retest in 2 days as recommended by the FDA. Child should wear a mask if age 2 years or older.</p> <p>STAY HOME IF child tests positive for COVID-19 (follow Isolation Instructions), if symptoms are accompanied by fever, or if child has significant nasal discharge.</p> <p>SEEK MEDICAL ATTENTION for severe symptoms, including a bad cough, difficulty breathing or trouble catching their breath, or wheezing when not controlled by medication (like albuterol).</p> <p>RETURN AFTER ISOLATION. If child tested positive for COVID-19, return after completing isolation per LACDPH isolation instructions.</p>



Guidance for Schools: Common Signs & Symptoms



Vomiting	<p>STAY HOME IF vomiting has occurred 2 or more times in 24 hours.</p> <p>RETURN WHEN vomiting has ended overnight, and child is able to hold down liquids and food. Recommend frequent handwashing.</p>
Diarrhea	<p>STAY HOME IF child's stool is likely to leak from the diaper, if they are unlikely to make it to the toilet in time (if potty trained), or if symptoms prevent them from participating meaningfully in routine activities.</p> <p>SEEK MEDICAL ATTENTION if the stool looks bloody or black.</p> <p>RETURN WHEN child has not had diarrhea for 24-48 hours or infectious cause has been ruled out. If child has been diagnosed with a specific infection, follow disease-specific guidance from public health or their healthcare provider.</p>
Sore Throat	<p>OK TO ATTEND WITH MILD SYMPTOMS. Child should wear a mask if age 2 years or older.</p> <p>SEEK MEDICAL ATTENTION for difficulty breathing or swallowing, or continuous drooling.</p> <p>RETURN WHEN IMPROVING. If an antibiotic is prescribed, take the first dose at least 12 hours before returning.</p>
Ear or eye irritation (including pink eye)	<p>OK TO ATTEND WITH MILD SYMPTOMS. Recommend frequent handwashing.</p> <p>STAY HOME IF there is significant discharge from the eye or ears, or if symptoms prevent child from participating meaningfully in routine activities.</p> <p>SEEK MEDICAL ATTENTION for difficulty seeing or hearing, an eye injury, or pain they cannot tolerate.</p>
Rash	<p>OK TO ATTEND WITH MILD SYMPTOMS.</p> <p>STAY HOME IF child has a fever or recent exposure to measles or chickenpox.</p> <p>SEEK MEDICAL ATTENTION if rash is worsening, causing pain, has blisters, or is not healing.</p> <p>RETURN WHEN IMPROVING or as guided by health care provider. In general, for conditions such as lice, impetigo, ringworm, scabies, and pinworms, your child may return as soon as they start appropriate treatment. Recommend covering rash if possible.</p>

Relevant Resources

[CDPH Guidance for K-12 Schools and Child Care, 2023-2024](#)

[CDPH Guidance When a Child Has Symptoms of Illness in School or Child Care](#)

[LA County COVID-19 Home Page](#)